

Degenerate Anisotropic Elliptic Problems and Magnetized Plasma Simulations

Stéphane Brull^{1,*}, Pierre Degond^{2,3} and Fabrice Deluzet^{2,3}

¹ *Institut de Mathématiques de Bordeaux UMR 5251, Equipe Mathématiques Appliquées de Bordeaux (MAB) Université Bordeaux 1 351, cours de la Libération-33405 TALENCE cedex, France.*

² *Université de Toulouse; UPS, INSA, UT1, UTM; Institut de Mathématiques de Toulouse; F-31062 Toulouse, France.*

³ *CNRS; Institut de Mathématiques de Toulouse UMR 5219, F-31062 Toulouse, France.*

Received 30 October 2010; Accepted (in revised version) 18 March 2011

Communicated by Kun Xu

Available online 5 September 2011

Abstract. This paper is devoted to the numerical approximation of a degenerate anisotropic elliptic problem. The numerical method is designed for arbitrary space-dependent anisotropy directions and does not require any specially adapted coordinate system. It is also designed to be equally accurate in the strongly and the mildly anisotropic cases. The method is applied to the Euler-Lorentz system, in the drift-fluid limit. This system provides a model for magnetized plasmas.

AMS subject classifications: 65N06, 65N12, 65M06, 65M12, 76W05, 76X05, 76N17

Key words: Anisotropic elliptic problem, asymptotic-preserving scheme, Lorentz force, large magnetic field, low-Mach number, drift-fluid limit.

1 Introduction

This paper discusses the numerical resolution of degenerate anisotropic elliptic problems of the form:

$$-(b \cdot \nabla)(\nabla \cdot (b\phi^\varepsilon)) + \varepsilon\phi^\varepsilon = f^\varepsilon, \quad \text{in } \Omega, \quad (1.1a)$$

$$(b \cdot \nu)\nabla \cdot (b\phi^\varepsilon) = 0, \quad \text{on } \partial\Omega, \quad (1.1b)$$

*Corresponding author. *Email addresses:* stephane.brull@math.u-bordeaux1.fr (S. Brull), pierre.degond@math.univ-toulouse.fr (P. Degond), fabrice.deluzet@math.univ-toulouse.fr (F. Deluzet)

where $\Omega \subset \mathbb{R}^2$ or \mathbb{R}^3 , f^ε is a given function, b is a normalized vector field defining the anisotropy direction and ε measures the strength of this anisotropy. In this expression ∇ and $\nabla \cdot$ are respectively the gradient and divergence operators. The unit outward normal at $x \in \partial\Omega$ is denoted by ν . In the context of plasmas, ε is related to the gyro period (i.e., the period of the gyration motion of the particles about the magnetic field lines) and the anisotropy direction b satisfies $b = B/|B|$ with the magnetic field B verifying $\nabla \cdot B = 0$. Eq. (1.1a) may also arise in other contexts, such as rapidly rotating flows, shell theory and may also be found when special types of semi-implicit time discretization of diffusion equations are used.

The elliptic equation is not in the usual divergence form due to an exchange between the gradient and divergence operators. However, the methodology would apply equally well to the operator $\nabla \cdot ((b \otimes b) \cdot \nabla \phi)$, up to some simple changes. The expression considered here is motivated by the application to the Euler-Lorentz system of plasmas. This application has already been considered in a previous study [13] but we introduce two important developments. First the present numerical method does not request the development of a special coordinate system adapted to b . In [13], b was assumed aligned with one coordinate direction. Second, the present paper considers Neumann boundary conditions instead of Dirichlet ones as in [13]. Although seemingly innocuous, this change brings in a considerable difficulty, linked with the degeneracy of the limit problem, as explained below.

A classical discretization of problem (1.1a), (1.1b) leads to an ill-conditioned linear system as $\varepsilon \rightarrow 0$. Indeed setting formally $\varepsilon = 0$ in (1.1a), (1.1b), we get:

$$-(b \cdot \nabla) \nabla \cdot (b\psi) = f^{(0)}, \quad \text{in } \Omega, \quad (1.2a)$$

$$(b \cdot \nu) \nabla \cdot (b\psi) = 0, \quad \text{on } \partial\Omega, \quad (1.2b)$$

with $f^{(0)} = \lim_{\varepsilon \rightarrow 0} f^\varepsilon$. The homogeneous system associated to (1.2a), (1.2b) admits an infinite number of solutions, namely all functions ψ satisfying $\nabla \cdot (b\psi) = 0$. This degeneracy results from the Neumann boundary conditions (1.2b) and would also occur if periodic boundary conditions were used. On the other hand, (1.2a) is not degenerate if supplemented with Dirichlet or Robin conditions, which was the case considered in [13]. A standard numerical approximation of (1.2a), (1.2b) generates a matrix whose condition number blows up as $\varepsilon \rightarrow 0$, leading to very time consuming and/or poorly accurate solution algorithms.

To bypass these limitations, we follow the idea introduced in [12] and use a decomposition of the solution in its average along the b -field lines and a fluctuation about this average. This decomposition ensures an accurate computation of the solution for all values of ε . In [12], this decomposition approach was developed for a uniform b and a coordinate system with one coordinate direction aligned with b . To extend this approach to arbitrary anisotropy fields b , a possible way is to use an adapted curvilinear coordinate system with one coordinate curve tangent to b . This is the route followed by [4], which proposes an extension of [12] in the context of ionospheric plasma physics, where the

anisotropy direction is known analytically (given by the earth dipolar magnetic field). The approach developed here is different and aims at a method which does not request the generation of special curvilinear coordinates. Indeed, in the general case, computing such coordinates can be complex and costly, especially for time-dependent problems where b evolves in time.

For this purpose, we solve a variational problem for each of the terms of the decomposition. The main difficulty lies in the discretization of the functional spaces in which each component of the solution is searched. In the present paper, this difficulty is solved by introducing two kinds of variational systems, one corresponding to a second-order elliptic problem (for the average) and one, to a fourth order system (for the fluctuation). An alternative to this method is proposed in [10]. It avoids the resolution of a fourth-order problem at the price of the introduction of Lagrange multipliers which lead to a larger system. In the present paper, we design a method which breaks the complexity of the problem in smaller pieces and requires less computer resources.

As an application of the method and a motivation for studying problem (1.2a), (1.2b), the drift-fluid limit of the isothermal Euler-Lorentz system is considered. These equations model the evolution of a magnetized plasma. In this case, the anisotropy direction is that of the magnetic field and the parameter ε is the reciprocal of the non dimensional cyclotron frequency. The drift-fluid limit $\varepsilon \rightarrow 0$ of the Euler-Lorentz system is singular because the momentum equation becomes degenerate. In this paper, we propose a scheme able to handle both the $\varepsilon \sim 1$ and $\varepsilon \ll 1$ regimes, giving rise to consistent approximations of both the Euler-Lorentz model and its drift-fluid limit, without any constraint on the space and time steps related to the possible small value of ε . Schemes having such properties are referred to as *Asymptotic-Preserving (AP)* schemes. These schemes are particularly efficient in situations in which part of the simulation domain is in the asymptotic regime and part of it is not. Indeed, in most practical cases, the parameter ε assumes a local value which may change from one location to the next or which may evolve with time.

The usual approach for dealing with such occurrences is through domain decomposition: the full Euler-Lorentz model is used in the region where $\varepsilon = \mathcal{O}(1)$ and the drift-fluid limit model is used where $\varepsilon \ll 1$. There are several drawbacks in using this approach. The first one is the choice of the position of the interface (or cross-talk region), which can influence the outcome of the simulation. If the interface evolves in time, an algorithm for interface motion has to be devised and some remeshing must be used to ensure compatibility between the mesh and the interface, which requires heavy code developments and can be quite CPU time consuming. Determining the right coupling strategy between the two models can also be quite challenging and the outcome of the numerical simulations may also depend on this choice. Because these questions do not have straightforward answers, domain decomposition strategies often lack robustness and reliability. Here, using the original model with an AP discretization method everywhere prevents from these artefacts and permits to use the same code everywhere for both regimes.

We conclude this introductory section by some bibliographical remarks. In magnetized plasma simulations, many works are based on the use of curvilinear coordinate

systems where one of the coordinate curves is tangent to the magnetic field (see e.g., [34], the gyro-kinetic and gyro-fluid developments [2, 19, 22, 24] and the many attempts for generating specialized coordinate systems [1, 5, 17, 18, 23, 26, 36]). The present work, together with [10] is one of the very few attempts to design numerical methods free of the use of special coordinate systems (see also [35]). The key idea behind this method is the concept of Asymptotic Preserving (AP) schemes as described above. AP-schemes have first been introduced by S. Jin [25] in the context of diffusive limits of transport models. They have recently found numerous applications to plasma physics in relation e.g., to quasineutrality [3, 9, 11, 14, 15] and strong magnetic fields [10, 12, 13] as well as to fluid-mechanical problems such as the small Mach-number limit of compressible fluids [16]. Other applications of AP-schemes can be found in [6–8, 20, 28, 29, 32]. Numerical methods for anisotropic problems have been extensively studied in the literature using numerous techniques such as domain decomposition techniques [21, 27], Multigrid methods, smoothers [31], the *hp*-finite element method [33]. However, these methods are based on a discretization of the anisotropic PDE as it is written. The method presented relies on a totally different concept, namely viewing the anisotropy as a singular perturbation and using Asymptotic-Preserving techniques.

For the readers convenience, we summarize the relations and differences with earlier work. In the present work as well as in [13], a fully degenerate elliptic equation of the form (1.1a) where the diffusion operator only acts in one direction is considered. However, [13] was concerned with a uniform anisotropy direction while the present work deals with the general case of a space-dependent anisotropy direction. Also, in [13], the boundary conditions were Dirichlet ones and the limit problem formally obtained by letting $\varepsilon \rightarrow 0$ in (1.1a) had still a unique solution. In the present case, the Neumann boundary conditions (1.1b) make the limit problem ill-posed and the $\mathcal{O}(\varepsilon)$ terms need to be considered to show that it is actually well-posed. This makes the numerics of the problem far more difficult. The works [4, 10, 12] deal with elliptic operators which are **not** degenerate until $\varepsilon = 0$ and finding the limit solution involves solving operators which are elliptic in the transverse direction. So, the setting is fairly different even if some methodologies can be and were borrowed. [12] introduces the decomposition of the solution that is also used in the present paper but is restricted to a uniform anisotropy direction. [4] is an extension to a space-dependent anisotropy direction which relies on the introduction of a special coordinate adapted to the anisotropy direction. Here, we want to avoid constructing such a coordinate system, which can be quite difficult and time-consuming in practice. [10] considers arbitrary coordinate systems but the constraints which apply to each component of the decomposition are dualized by means of Lagrange multipliers, which greatly increases the number of unknowns and the computer time. In the present work, these constraints are expressed through some functional properties, that take advantage of the fully degenerate form of the elliptic problem to be solved. For the sake of the efficiency of the numerical exploration, the numerical experiments have been restricted to two space dimensions. Fully three-dimensional simulations are in progress and will be reported in future work.

This paper is organized as follows. In Section 2 the solution methodology for the degenerate anisotropic elliptic problem (1.1a), (1.1b) is detailed and the variational method avoiding the use of adapted coordinates is presented. Section 3 is devoted to the discretization strategy. The numerical method for the anisotropic elliptic problem is validated in Section 4. In Section 5 the drift-fluid limit of the isothermal Euler-Lorentz system is introduced. The AP-scheme is derived, giving rise to the anisotropic elliptic problem (1.1a), (1.1b). Finally, numerical comparisons of the classical and AP schemes for the Euler-Lorentz system are given in Section 6. A conclusion is drawn in Section 7.

2 A decomposition method for degenerate anisotropic elliptic problems

We first present the methodology in the simpler case of a uniform b -field. The method will then be extended to an arbitrary b -field.

2.1 Overview of the method in the uniform b -field case

A two dimensional configuration is considered in this section, with the position variable (x,y) belonging to a square domain $(x,y) \in \Omega = [0,1] \times [0,1] \subset \mathbb{R}^2$. The b field is assumed uniform, equal to the unit vector pointing in the y direction. In this case, the singular perturbation problem (1.1a), (1.1b) reads:

$$\varepsilon \phi^\varepsilon(x,y) - \frac{\partial^2}{\partial y^2} \phi^\varepsilon(x,y) = f^\varepsilon(x,y), \quad \text{in } [0,1] \times [0,1], \quad (2.1a)$$

$$\frac{\partial}{\partial y} \phi^\varepsilon(x,y) = 0, \quad \text{for } y=0 \text{ or } y=1. \quad (2.1b)$$

We assume that:

$$\lim_{\varepsilon \rightarrow 0} \left(\frac{1}{\varepsilon} \int_0^1 f^\varepsilon(x,y) dy \right) \text{ exists and is finite, } \forall x \in [0,1]. \quad (2.2)$$

This framework is similar to [12]. Here, we recall the bases of the methodology. The problem is well posed for all $\varepsilon > 0$ but a standard discretization may lead to ill-conditioned matrices when $\varepsilon \ll 1$. Indeed if ε is formally set to zero, we get the following degenerate problem

$$-\frac{\partial^2}{\partial y^2} \psi(x,y) = f^{(0)}(x,y), \quad \text{in } [0,1] \times [0,1], \quad (2.3a)$$

$$\frac{\partial}{\partial y} \psi(x,y) = 0, \quad \text{for } y=0 \text{ or } y=1, \quad (2.3b)$$

assuming that f^ε has the following expansion $f^\varepsilon = f^{(0)} + \varepsilon f^{(1)} + o(\varepsilon)$. This system admits a solution under the compatibility condition $\int_0^1 f^{(0)}(x,y) dy = 0$ for all $x \in [0,1]$, which is

satisfied thanks to hypothesis (2.2). However the solution is not unique. Indeed, if ψ verifies (2.3) then $\psi + \zeta$ is also a solution for all functions $\zeta = \zeta(x)$ which depend on the x -coordinate only.

On the other hand, the limit $\phi^{(0)} = \lim_{\varepsilon \rightarrow 0} \phi^\varepsilon$ is unique. Indeed, it is easy to see that the solution $\tilde{\psi}$ of (2.3) such that $\int_0^1 \tilde{\psi}(x, y) dy = 0$ for all $x \in [0, 1]$ is unique. Since $\phi^{(0)}$ is a particular solution of (2.3), it can be written

$$\phi^{(0)} = \tilde{\psi} + \zeta(x). \quad (2.4)$$

In order to determine ζ , we integrate (2.1a) with respect to y and get

$$\int_0^1 \phi^\varepsilon(x, y) dy = \frac{1}{\varepsilon} \int_0^1 f^\varepsilon(x, y) dy. \quad (2.5)$$

Taking the limit $\varepsilon \rightarrow 0$ in this equation and inserting (2.4), we get $\zeta(x) = \int_0^1 f^{(1)}(x, y) dy$, which determines $\phi^{(0)}$ uniquely.

Now, if a standard numerical method is applied to (2.1a), (2.1b), the resulting matrix will be close, when $\varepsilon \ll 1$, to the singular matrix obtained from the discretization of (2.3). Therefore, its condition number will blow up as $\varepsilon \rightarrow 0$, resulting in either low accuracy, or high computational cost. To overcome this problem, we decompose ϕ^ε according to

$$\phi^\varepsilon = p^\varepsilon + q^\varepsilon, \quad p^\varepsilon(x) = \int_0^1 \phi^\varepsilon(x, y) dy, \quad (2.6)$$

i.e., p^ε is the average of ϕ^ε along straight lines parallel to b and q^ε is the fluctuation of the solution with respect to this average. p^ε and q^ε satisfy:

$$\frac{\partial p^\varepsilon}{\partial y}(x, y) = 0, \quad \forall (x, y) \in \Omega, \quad (2.7a)$$

$$\int_0^1 q^\varepsilon(x, y) dy = 0, \quad \forall x \in [0, 1]. \quad (2.7b)$$

They are orthogonal for the scalar product of L^2 , i.e., $\int_\Omega p^\varepsilon q^\varepsilon dx dy = 0$.

Inserting this decomposition into (2.5) yields

$$p^\varepsilon(x) = \frac{1}{\varepsilon} \int_0^1 f^\varepsilon(x, y) dy, \quad \forall x \in [0, 1]. \quad (2.8)$$

Moreover, p^ε satisfies

$$\lim_{\varepsilon \rightarrow 0} p^\varepsilon(x) = \lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} \int_0^1 f^\varepsilon(x, y) dy = \int_0^1 f^{(1)}(x, y) dy = \zeta(x),$$

where ζ is defined by (2.4). Now, q^ε is the solution of the following problem:

$$-\frac{\partial^2}{\partial y^2}q^\varepsilon(x,y) + \varepsilon q^\varepsilon(x,y) = \zeta^\varepsilon(x,y), \quad \forall (x,y) \in [0,1] \times [0,1], \quad (2.9a)$$

$$\int_0^1 q^\varepsilon(x,y)dy = 0, \quad \text{for } x \in [0,1], \quad (2.9b)$$

$$\frac{\partial}{\partial y}q^\varepsilon(x,y) = 0, \quad \text{for } y=0 \text{ or } y=1, \quad (2.9c)$$

where

$$\zeta^\varepsilon = f^\varepsilon - \int_0^1 f^\varepsilon dy = f^\varepsilon - \varepsilon p^\varepsilon$$

is the projection of f^ε on the space of functions satisfying (2.7b). Compared to (2.1a), (2.1b), system (2.9a)-(2.9c) involves the additional condition (2.9b). This condition is important: it makes the system uniformly well-posed when $\varepsilon \rightarrow 0$. Additionally, the limit system is

$$-\frac{\partial^2}{\partial y^2}q^{(0)}(x,y) = f^{(0)}, \quad \forall (x,y) \in [0,1] \times [0,1], \quad (2.10a)$$

$$\int_0^1 q^{(0)}(x,y)dy = 0, \quad \text{for } x \in [0,1], \quad (2.10b)$$

$$\frac{\partial}{\partial y}q^{(0)}(x,y) = 0, \quad \text{for } y=0 \text{ or } y=1, \quad (2.10c)$$

and has a unique solution equal to $\tilde{\psi}$. Consequently, as $\varepsilon \rightarrow 0$

$$\phi^\varepsilon = p^\varepsilon + q^\varepsilon \rightarrow \zeta + \tilde{\psi} = \phi^{(0)}.$$

Therefore, the proposed decomposition leads to two uniformly well-posed problems when $\varepsilon \rightarrow 0$, which allows to reconstruct the limit solution $\phi^{(0)}$ of the original problem.

The numerical approximations of conditions (2.8) or (2.9b) is delicate if the mesh is not aligned with the y coordinate axis. In order to overcome this problem, a weak formulation is introduced. Define $V = H^1(0,1)$, $K = \{v \in V \mid \partial_y v = 0\}$. Then, ϕ^ε is the solution of the variational formulation: find $\phi^\varepsilon \in V$ such that

$$\int_\Omega \frac{\partial \phi^\varepsilon}{\partial y} \frac{\partial \psi}{\partial y} dx dy + \varepsilon \int_\Omega \phi^\varepsilon \psi dx dy = \int_\Omega f^\varepsilon \psi dx dy, \quad \forall \psi \in V. \quad (2.11)$$

Let K^\perp be the orthogonal space to K in $L^2(0,1)$. Now, the decomposition (2.6), corresponds to the decomposition of ϕ^ε on K and K^\perp . Indeed, it is easily checked that $p^\varepsilon \in K$ and $q^\varepsilon \in K^\perp$ and they are orthogonal, as already noticed. Now, inserting $\psi \in K$ in (2.11), we get that p^ε is the solution of: find $p^\varepsilon \in K$ such that

$$\int_\Omega \left(p^\varepsilon - \frac{1}{\varepsilon} f^\varepsilon \right) \psi dx dy = 0, \quad \forall \psi \in K, \quad (2.12)$$

which means that p^ε is the orthogonal projection of $\varepsilon^{-1}f^\varepsilon$ onto K . Now, inserting $\psi \in K^\perp$ in (2.11) leads to: find $q^\varepsilon \in K^\perp$ such that

$$\int_{\Omega} \frac{\partial q^\varepsilon}{\partial y} \frac{\partial \psi}{\partial y} dx dy + \varepsilon \int_{\Omega} q^\varepsilon \psi dx dy = \int_{\Omega} (f^\varepsilon - \varepsilon p^\varepsilon) \psi dx dy, \quad \forall \psi \in K^\perp, \quad (2.13)$$

which is the variational formulation of (2.9a)-(2.9c).

The use of these variational formulations allows for the discretization of (2.1a), (2.1b) on arbitrary meshes compared to the anisotropy direction. This is an important advantage over the strong formulations (2.8) or (2.9a)-(2.9c). These formulations are now generalized to arbitrary anisotropy fields b in the next section.

2.2 Presentation of the method for a general anisotropy field

In this section, we introduce the functional setting for the variational formulation of the anisotropic elliptic problem in the variable b case. This variational method is the cornerstone of the strategy presented in this paper and avoids the use of curvilinear coordinates adapted to b .

2.2.1 Preliminaries

This subsection is devoted to the resolution of degenerate elliptic problems (1.1a), (1.1b) for general anisotropy fields b . we first introduce the space

$$\begin{aligned} \mathcal{V} &= \{ \phi \in L^2(\Omega) / \nabla \cdot (b\phi) \in L^2(\Omega) \}, & K &= \{ \phi \in \mathcal{V} / \nabla \cdot (b\phi) = 0 \text{ on } \Omega \}, \\ \mathcal{W} &= \{ h \in L^2(\Omega) / (b \cdot \nabla)h \in L^2(\Omega) \}, & \mathcal{W}_0 &= \{ h \in \mathcal{W} / (b \cdot \nu)h = 0 \text{ on } \partial\Omega \}. \end{aligned}$$

The projection of a function on K is the generalization of the average operation (2.8), while the projection on K^\perp corresponds to computing its fluctuation. The space \mathcal{W}_0 is used to characterize K^\perp . The projections on K and K^\perp are well-defined thanks to the:

Theorem 2.1. *We have the following properties*

- 1) K is closed in $L^2(\Omega)$.
- 2) \mathcal{W}_0 equipped with the norm $\|h\|_{\mathcal{W}_0} = \|(b \cdot \nabla)h\|_{L^2(\Omega)}$ is a Hilbert space and $(b \cdot \nabla)\mathcal{W}_0$ is a closed space of $L^2(\Omega)$.
- 3) $K^\perp = (b \cdot \nabla)\mathcal{W}_0$.

Proof. 1) Let $\phi_n \in \mathcal{V}$ such that $\phi_n \rightarrow \phi$ in $L^2(\Omega)$. Then, $\phi_n \rightarrow \phi$ in the distributional sense and the operation $\phi \rightarrow \nabla \cdot (b\phi)$ is continuous for the topology of distributions. Therefore, $\nabla \cdot (b\phi) = 0$, which shows that $\phi \in \mathcal{V}$.

2) \mathcal{W}_0 is a Hilbert space for the norm $\|h\| = \|h\|_{L^2(\Omega)} + \|(b \cdot \nabla)h\|_{L^2(\Omega)}$. According to the Poincaré inequality, the norms $\| \cdot \|$ and $\| \cdot \|_{\mathcal{W}_0}$ are equivalent. The closedness of \mathcal{W}_0 for the L^2 topology follows from 3).

3) The inclusion $(b \cdot \nabla) \mathcal{W}_0 \subseteq K^\perp$ is obvious. We sketch the proof of the converse inclusion and leave the details to the reader. We make the hypothesis that all b -field lines are either tangent to a non-zero measure set of $\partial\Omega$ or intersect $\partial\Omega$ at two points x_- and x_+ such that $\pm(b \cdot \nu)(x_\pm) > 0$. The points x_- and x_+ are called the conjugate points of the b -field line and are respectively the incoming and outgoing points of this field line to the domain. These assumptions can certainly be weakened at the expense of technical difficulties which are outside the scope of this paper. Let $\psi \in K^\perp$. By taking the primitive of ψ along the b -field lines, there exists $\phi \in \mathcal{W}$ such that $\psi = (b \cdot \nabla)\phi$. We can additionally impose that $\phi = 0$ on $\partial\Omega_-$, where $\partial\Omega_\pm = \{x \in \partial\Omega \mid \pm(b \cdot \nu)(x) > 0\}$. Let $\theta \in K$. We have

$$0 = \int_\Omega \psi \theta dx = \int_\Omega (b \cdot \nabla)\phi \theta dx = \int_{\partial\Omega} (b \cdot \nu)\phi \theta dS(x), \tag{2.14}$$

where $dS(x)$ is the superficial measure on $\partial\Omega$. Since, $\theta \in K$ its values at conjugate points are related by a linear relation. In particular, they can be taken simultaneously non-zero. Then, since the values of ϕ on $\partial\Omega_-$ vanish, (2.14) implies that the values of ϕ on $\partial\Omega_+$ vanish as well. Consequently, $(b \cdot \nu)\phi = 0$ on $\partial\Omega$, which shows that $\phi \in \mathcal{W}_0$. This proves the result. \square

Therefore, we can decompose ϕ^ε uniquely as

$$\phi^\varepsilon = p^\varepsilon + q^\varepsilon, \quad p^\varepsilon \in K, \quad q^\varepsilon \in K^\perp, \tag{2.15}$$

and state problem (1.1a), (1.1b) as

$$-(b \cdot \nabla)(\nabla \cdot (bq^\varepsilon)) + \varepsilon(p^\varepsilon + q^\varepsilon) = f^\varepsilon, \quad \text{in } \Omega, \tag{2.16a}$$

$$(b \cdot \nu)\nabla \cdot (bq^\varepsilon) = 0, \quad \text{in } \partial\Omega, \tag{2.16b}$$

$$p^\varepsilon \in K \text{ and } q^\varepsilon \in K^\perp. \tag{2.16c}$$

Next, we introduce the variational approach. We multiply (2.16a) by a test function $\psi \in \mathcal{V}$ and integrate it on Ω . Using a Green formula together with the boundary condition (2.16b), we find that

$$\int_\Omega \nabla \cdot (bq^\varepsilon) \nabla \cdot (b\psi) dx + \varepsilon \int_\Omega (p^\varepsilon + q^\varepsilon) \psi dx = \int_\Omega f^\varepsilon \psi dx. \tag{2.17}$$

Now, the aim is to decompose problem (2.17) into a problem for p^ε and a problem for q^ε . For this purpose, in the following section, the test function ψ is chosen successively in K and in K^\perp .

2.2.2 Equation for $p^\varepsilon \in K$ and $q^\varepsilon \in K^\perp$

The following proposition is devoted to the determination of for $p^\varepsilon \in K$ and $q^\varepsilon \in K^\perp$.

Proposition 2.1. Let p^ε and q^ε be given by (2.15) where ϕ^ε is the solution of problem (1.1a), (1.1b). Then:

(i) p^ε is given by

$$p^\varepsilon = \frac{1}{\varepsilon}(f^\varepsilon + b \cdot \nabla g^\varepsilon), \quad \text{in } \Omega, \quad (2.18)$$

where g^ε satisfies the problem

$$-\nabla \cdot ((b \otimes b) \nabla g^\varepsilon) = \nabla \cdot (f^\varepsilon b), \quad \text{in } \Omega, \quad (2.19a)$$

$$(b \cdot \nu) g^\varepsilon = 0, \quad \text{on } \partial\Omega, \quad (2.19b)$$

or, in variational form, find $g^\varepsilon \in \mathcal{W}_0$ such that

$$\int_{\Omega} (b \cdot \nabla g^\varepsilon)(b \cdot \nabla \theta) = \int_{\Omega} f^\varepsilon b \cdot \nabla \theta dx, \quad \forall \theta \in \mathcal{W}_0. \quad (2.20)$$

(ii) q^ε is given by

$$q^\varepsilon = b \cdot \nabla h^\varepsilon, \quad (2.21)$$

where h^ε satisfies the following fourth-order problem:

$$-\nabla \cdot [(b \otimes b) \nabla (\nabla \cdot (b \otimes b) \nabla h^\varepsilon)] + \varepsilon \nabla \cdot ((b \otimes b) \nabla h^\varepsilon) = \nabla \cdot (b f^\varepsilon), \quad \text{in } \Omega, \quad (2.22a)$$

$$(b \cdot \nu) \nabla \cdot ((b \otimes b) \nabla h^\varepsilon) = 0, \quad \text{on } \partial\Omega, \quad (2.22b)$$

$$(b \cdot \nu) h^\varepsilon = 0, \quad \text{on } \partial\Omega, \quad (2.22c)$$

or, in variational form, find $h^\varepsilon \in \mathcal{W}_0$ such that

$$\int_{\Omega} \nabla \cdot ((b \otimes b) \nabla h^\varepsilon) \nabla \cdot ((b \otimes b) \nabla \theta) dx + \varepsilon \int_{\Omega} (b \cdot \nabla h^\varepsilon)(b \cdot \nabla \theta) dx = \int_{\Omega} f^\varepsilon (b \cdot \nabla \theta) dx. \quad (2.23)$$

Proof. Choosing $\psi = r \in K$ in (2.17), we obtain the problem, find $p^\varepsilon \in K$ such that

$$\int_{\Omega} (\varepsilon p^\varepsilon - f^\varepsilon) r dx = 0, \quad \forall r \in K. \quad (2.24)$$

This problem admits a solution in K which is uniformly bounded in $L^2(\Omega)$ as $\varepsilon \rightarrow 0$ under the compatibility condition

$$\lim_{\varepsilon \rightarrow 0} \left(\frac{1}{\varepsilon} \int_{\Omega} f^\varepsilon r dx \right) \text{ exists and is finite, } \forall r \in K. \quad (2.25)$$

Assuming that f^ε has the following decomposition

$$f^\varepsilon = f^{(0)} + \varepsilon f^{(1)} + o(\varepsilon)$$

in $L^2(\Omega)$, this condition implies that $f^{(0)} \in K^\perp$. Next, since $\varepsilon p^\varepsilon - f^\varepsilon \in K^\perp$, according to Theorem 2.1, there exists $g^\varepsilon \in \mathcal{W}_0$ such that

$$\varepsilon p^\varepsilon - f^\varepsilon = (b \cdot \nabla) g^\varepsilon. \quad (2.26)$$

Taking the product with b and the divergence of the result, we obtain (2.19a) and (2.19b).

Taking $\psi = s \in K^\perp$ in (2.17) gives:

$$\int_{\Omega} \nabla \cdot (b q^\varepsilon) \nabla \cdot (b s) dx + \varepsilon \int_{\Omega} q^\varepsilon s dx = \int_{\Omega} f^\varepsilon s dx. \quad (2.27)$$

But since q^ε and $s \in K^\perp$, Theorem 2.1 implies that there exists h^ε and $\theta \in \mathcal{W}_0$ such that $q^\varepsilon = b \cdot \nabla h^\varepsilon$ and $s = b \cdot \nabla \theta$. \square

Remark 2.1. In [10], the characterization of K^\perp as $(b \cdot \nabla) \mathcal{W}_0$ is not used. Instead, the constraint that $q \in K^\perp$ is taken into account through a mixed formulation. The number of unknowns and the size of the problem are therefore larger in [10] than in the present work. In practice, the resolution of the fourth order problem (2.22a), (2.22b), (2.22c) can be reduced by solving two second-order problem, as shown below. Therefore, the introduction of a fourth order problem does not bring specific difficulties.

2.2.3 Extension to non-homogeneous Neumann boundary conditions

The application targeted in this paper and detailed in Section 5, requires the handling of non-homogeneous Neumann boundary conditions. In this subsection ϕ^ε is solution to the following inhomogeneous Neumann problem:

$$\varepsilon \phi^\varepsilon - (b \cdot \nabla) (\nabla \cdot (b \phi^\varepsilon)) = b \cdot \nabla \kappa + f_2^\varepsilon, \quad \text{on } \Omega, \quad (2.28a)$$

$$(b \cdot \nu) \nabla \cdot (b \phi^\varepsilon) = -(b \cdot \nu) \kappa, \quad \text{on } \partial \Omega, \quad (2.28b)$$

where κ is a given function in \mathcal{W} . We denote by $f_1 = b \cdot \nabla \kappa$ and by $f^\varepsilon = f_1 + f_2^\varepsilon$.

Using the same decomposition (2.15) as before, we find that p^ε satisfies (2.18) and g^ε is the solution of (2.19a), (2.19b) or (2.20) with f^ε replaced by f_2^ε (and satisfying (2.25)). Similarly, q^ε satisfies (2.21) and h^ε is the solution of (2.22a), (2.22b), (2.22c), or of (2.23) with "0" at the right-hand side of (2.22b) replaced by $(b \cdot \nu) \kappa$, the other terms being unchanged. The details are left to the reader.

3 Space discretization


In this section, we investigate the finite-volume discretization of the two elliptic problems arising in Proposition 2.1. The plan is to use the resulting degenerate elliptic solver as a building block for the AP-discretization of the Euler-Lorentz model (see Section 5). For systems of conservation laws like the Euler-Lorentz model, finite-volume methods are a natural choice given the immense available scientific literature (see [30] and references

therein). We also restrict ourselves to structured meshes, because our primary objective is to test the AP methodology in a simple setting. Additionally, the applications we are targeting (such as tokamaks), as well as many other applications can be developed using a logically structured mesh. In future work, the extension of the method to unstructured meshes will be investigated, particularly in view of adaptive mesh refinement. We note that the finite-volume setting on structured meshes can be equivalently formulated into a finite-difference setting. However, we prefer the finite-volume setting in relation to the final discretization of the Euler-Lorentz model as stated before.

The domain is decomposed into a family \mathcal{R} of rectangles $M_{i-1/2,j-1/2} = [x_{i-1}, x_i] \times [y_{j-1}, y_j]$ with $x_i = i\Delta x$ and $y_j = j\Delta y$. We look for a piecewise constant approximation $p_{\mathcal{R}}^\varepsilon$ of p^ε on each $M_{i-1/2,j-1/2}$ and denote by $p_{i-1/2,j-1/2}$ its constant value on this rectangle. The function g^ε is approximated by a constant function on a dual mesh \mathcal{D} , consisting of rectangles $\mathcal{D}_{i,j} = [x_{i-1/2}, x_{i+1/2}] \times [y_{j-1/2}, y_{j+1/2}]$, where $x_{i-1/2} = (i-1/2)\Delta x$, $y_{i-1/2} = (i-1/2)\Delta y$. Then g^ε is approximated by a piecewise constant function $g_{\mathcal{D}}^\varepsilon$ with its constant values denoted by $g_{i,j}^\varepsilon$. We approximate (2.18) by

$$p_{i-\frac{1}{2},j-\frac{1}{2}} = \frac{1}{\varepsilon} \left(f^\varepsilon(x_{i-\frac{1}{2}}, y_{j-\frac{1}{2}}) + b(x_{i-\frac{1}{2}}, y_{j-\frac{1}{2}}) \cdot (\nabla g^\varepsilon)_{i-\frac{1}{2},j-\frac{1}{2}} \right),$$

where $b(x_{i-1/2}, y_{j-1/2}) \cdot (\nabla g^\varepsilon)_{i-1/2, j-1/2}$ is given in (3.1). We now define approximations $(b \cdot \nabla)_{\text{app}}$ and $\nabla \cdot (b \cdot \Phi)_{\text{app}}$ of operators $\Psi \mapsto (b \cdot \nabla \Psi)$ and $\Phi \mapsto \nabla \cdot (b \Phi)$ such that they are discrete dual operators to each other. For this purpose, we define $L_{\mathcal{R}}$ and $L_{\mathcal{D}}$ the space of piecewise constant functions on meshes of types \mathcal{R} and \mathcal{D} respectively.



Definition 3.1. The operator $(b \cdot \nabla)_{\text{app}}: L_{\mathcal{D}} \rightarrow L_{\mathcal{R}}$ is defined by

$$\begin{aligned} ((b \cdot \nabla \Psi)_{\text{app}})_{i-\frac{1}{2},j-\frac{1}{2}} = b(x_{i-\frac{1}{2}}, y_{j-\frac{1}{2}}) \cdot \left(\left(\frac{\Psi_{i,j} - \Psi_{i-1,j}}{2\Delta x} + \frac{\Psi_{i,j-1} - \Psi_{i-1,j-1}}{2\Delta x} \right), \right. \\ \left. \left(\frac{\Psi_{i,j} - \Psi_{i,j-1}}{2\Delta y} + \frac{\Psi_{i-1,j} - \Psi_{i-1,j-1}}{2\Delta y} \right) \right). \end{aligned} \quad (3.1)$$

The operator $\nabla \cdot (\cdot b)_{\text{app}}: L_{\mathcal{R}} \rightarrow L_{\mathcal{D}}$ is defined by

$$\begin{aligned} (\nabla \cdot (b\Phi)_{\text{app}})_{i,j} = & \left(\frac{1}{2\Delta x} b_x(x_{i+\frac{1}{2}}, y_{j-\frac{1}{2}}) - \frac{1}{2\Delta y} b_y(x_{i+\frac{1}{2}}, y_{j-\frac{1}{2}}) \right) \Phi_{i+\frac{1}{2}, j-\frac{1}{2}} \\ & + \left(\frac{1}{2\Delta x} b_x(x_{i+\frac{1}{2}}, y_{j+\frac{1}{2}}) + \frac{1}{2\Delta y} b_y(x_{i+\frac{1}{2}}, y_{j+\frac{1}{2}}) \right) \Phi_{i+\frac{1}{2}, j+\frac{1}{2}} \\ & - \left(\frac{1}{2\Delta x} b_x(x_{i-\frac{1}{2}}, y_{j-\frac{1}{2}}) + \frac{1}{2\Delta y} b_y(x_{i-\frac{1}{2}}, y_{j-\frac{1}{2}}) \right) \Phi_{i-\frac{1}{2}, j-\frac{1}{2}} \\ & - \left(\frac{1}{2\Delta x} b_x(x_{i-\frac{1}{2}}, y_{j+\frac{1}{2}}) - \frac{1}{2\Delta y} b_y(x_{i-\frac{1}{2}}, y_{j+\frac{1}{2}}) \right) \Phi_{i-\frac{1}{2}, j+\frac{1}{2}}. \end{aligned} \quad (3.2)$$

Proposition 3.1. $(b \cdot \nabla)_{\text{app}}$ and $\nabla \cdot (b \cdot)_{\text{app}}$ are adjoint operators to each other.

Proof. Easy and left to the reader, thanks to a discrete Green formula. \square

Next, we define $(\nabla \cdot ((b \otimes b) \cdot \nabla))_{\text{app}}$ by the composition of the two operators $(b \cdot \nabla)_{\text{app}}$ and $\nabla \cdot (b \cdot)_{\text{app}}$:

Definition 3.2. We define:

$$\left(\nabla \cdot (b \otimes b \cdot \nabla) \right)_{\text{app}} = (\nabla \cdot (\cdot b))_{\text{app}} \circ (b \cdot \nabla)_{\text{app}}, \quad (3.3)$$

where \circ is the composition operation.

Finally, the approximation of problem (2.19a), (2.19b) is by solving the discrete problem for the piecewise constant function g on \mathcal{D} :

$$\left(\nabla \cdot (b \otimes b \cdot \nabla) \right)_{\text{app}} g = (\nabla \cdot (bf))_{\text{app}}, \quad (3.4)$$

together with Dirichlet boundary conditions on g , where f is a piecewise constant function on \mathcal{R} .

Now, problem (2.22a), (2.22b), (2.22c) for q^ε can be decomposed in two decoupled second-order elliptic problems of the type (2.19a), (2.19b) and can be solved by a similar method. Indeed by setting $u = -\nabla \cdot ((b \otimes b) \nabla h)$, we get that (2.22a), (2.22b), (2.22c) is equivalent to the following two elliptic problems:

$$\nabla \cdot ((b \otimes b) \nabla u) - \varepsilon u = \nabla \cdot (bf), \quad \text{in } \Omega, \quad (3.5a)$$

$$(b \cdot \nu)u = 0, \quad \text{on } \partial\Omega \quad (3.5b)$$

and

$$-\nabla \cdot ((b \otimes b) \nabla h) = u, \quad \text{in } \Omega, \quad (3.5c)$$

$$(b \cdot \nu)h = 0, \quad \text{on } \partial\Omega. \quad (3.5d)$$

To summarize, the resolution of problem (1.1a), (1.1b) reduces to three independent resolutions of problems similar to (3.4).

4 Numerical results for the elliptic problem

4.1 Introduction

In this section the efficiency of the numerical method introduced in Sections 2 and 3 for the singular perturbation problem (1.1a), (1.1b) is investigated through numerical experiments. These experiments are carried out on a two dimensional uniform Cartesian mesh. Two sets of test cases are presented. In the first one, the anisotropy, or magnetic field, is oblique, which means that it is assumed uniform in space, but not necessarily aligned with any coordinate axis. In the second set, the field direction is non uniform. In both cases, the strength of the anisotropy is assumed uniform and is given by the value of ε . An analytical solution ϕ_a is constructed for the singular perturbation problem (1.1a), (1.1b) and is compared with its approximation ϕ^h computed on the mesh. For the test cases, the following L^1 , L^2 and L^∞ norms are used to estimate the errors between the numerical approximation ϕ^h and the analytical solution $\tilde{\phi}_a$:

$$e_1 = \frac{\|\tilde{\phi}_a - \phi^h\|_{L^1}}{\|\tilde{\phi}_a\|_{L^1}} = \frac{\sum_{i,j} |\phi_a(x_i, y_j) - \phi^h(i, j)|}{\sum_{i,j} |\phi_a(x_i, y_j)|}, \quad (4.1a)$$

$$e_2 = \frac{\|\tilde{\phi}_a - \phi^h\|_{L^2}}{\|\tilde{\phi}_a\|_{L^2}} = \frac{(\sum_{i,j} |\phi_a(x_i, y_j) - \phi_{i,j}^h|^2)^{\frac{1}{2}}}{(\sum_{i,j} |\phi_a(x_i, y_j)|^2)^{\frac{1}{2}}}, \quad (4.1b)$$

$$e_\infty = \frac{\|\tilde{\phi}_a - \phi^h\|_{L^\infty}}{\|\tilde{\phi}_a\|_{L^\infty}} = \frac{\max_{i,j} |\phi_a(x_i, y_j) - \phi_{i,j}^h|}{\max_{i,j} |\phi_a(x_i, y_j)|}. \quad (4.1c)$$

4.2 Numerical results for an oblique magnetic field

4.2.1 Introduction and test case settings

For these numerical experiments the simulation domain is the square $\Omega = [0,1] \times [0,1]$. The magnetic field is defined by $B = (\sin\alpha, \cos\alpha, 0)$, with α the angle of the b -field with the x -axis ranging from 0 to $\pi/2$. In order to validate the numerical method an analytical solution denoted ϕ_a for problem (1.1a), (1.1b) is constructed. It is written

$$\phi_a(x, y) = \sin(x \sin(\alpha) - y \cos(\alpha)) + b \cdot \nabla H(x, y), \quad (4.2a)$$

$$f_a^\varepsilon(x, y) = -b \cdot \nabla (\nabla \cdot ((b \otimes b) \nabla H(x, y))) + \varepsilon (\sin(x \sin(\alpha) - y \cos(\alpha)) + b \cdot \nabla H(x, y)), \quad (4.2b)$$

$$H(x, y) = ((x-1)(y-1)xy)^3. \quad (4.2c)$$

The function ϕ_a is the solution of problem (1.1a), (1.1b) with the right-hand side f_a^ε . ϕ_a presents itself as decomposed into p^ε (first terms) and q^ε (second term). Note also that f_a^ε can be decomposed as $f_a^\varepsilon = f_a^{(0)} + \varepsilon f_a^{(1)}$ with $f_a^{(0)} = -b \cdot \nabla h$ and $h = \nabla \cdot ((b \otimes b) \nabla H(x, y))$. The function h verifies homogeneous Dirichlet boundary conditions on the domain boundaries, which implies, according to Theorem 2.1, that $f_a^{(0)} \in K^\perp$ and the compatibility con-

dition (2.25) is satisfied. However, for the simulations carried out below, the construction of the right-hand side f_a^ε is performed using the discrete operators $(b \cdot \nabla)_{\text{app}}$ and $\nabla \cdot (\cdot b)_{\text{app}}$ in order to ensure that the compatibility condition (2.25) is satisfied by the discrete operators, namely $f_a^{(0)} \in K_{\text{app}}^\perp$, where

$$K_{\text{app}} = \{ \phi / \nabla \cdot (b\phi)_{\text{app}} = 0 \}, \quad K_{\text{app}}^\perp = (b \cdot \nabla)_{\text{app}}(\mathcal{W}_0).$$

4.2.2 Homogeneous Neumann boundary conditions

This simulation is run with $\alpha = \pi/3$. On Fig. 1, we represent the relative errors as functions of the mesh sizes for different values of ε ranging from 10^{-3} to 10^{-9} . The curves of Fig. 1 are plotted using logarithmic decimal scales. We observe a linear decrease of the errors with vanishing mesh sizes, with a slope equal to 2, which proves that the global scheme is second order accurate. More importantly, we observe from Figs. 1(a) and 1(b), that the precision remains the same while ε is decreased by three orders of magnitude. However, for the more refined grids using the smallest value of ε of this simulation set (10^{-9} , see Fig. 1(c)), a slight degradation of the convergence is observed for small mesh sizes.

This slight degradation can be explained. Indeed, p^ε is given by a stiff problem, since $\varepsilon p^\varepsilon$ is obtained as the difference of two quantities scaling as $\varepsilon^0 = \mathcal{O}(1)$ (see (2.18), (2.19a)). To investigate the influence of ε on the accuracy of the approximation of p^ε , the L^∞ norm of the relative error made on p^ε and on $\nabla \cdot (bp^\varepsilon)$ as functions of ε are plotted on Fig. 2.

Fig. 2(a) shows a linear behavior of $\nabla \cdot (bp^\varepsilon)$ with vanishing ε (in log scale). To explain this feature, we note that the discretization of the second order operator in (2.19a) provides a computation of $\varepsilon(\nabla \cdot (\cdot b))_{\text{app}}(p^\varepsilon)$ with the precision of the linear system solver used for the computation of g^ε , which is limited by round-off errors. This error is amplified after multiplication by the factor $1/\varepsilon$. This analysis still holds for the accuracy of p^ε as a function of ε represented on Fig. 2(b) with slight differences. For the largest values of ε , we observe a plateau (red dashed line) explained by the discretization error of the discrete operators. The space discretization introduced here is second order accurate, i.e., is $\mathcal{O}(h^2)$ where $h = \max(\Delta x, \Delta y)$. Since the right-hand side is well prepared this error only applies to the $\varepsilon f^{(1)}$ part of f^ε and is then proportional to $\varepsilon \mathcal{O}(h^2)$ in $b \cdot \nabla g^\varepsilon$, giving rise to a $\mathcal{O}(h^2)$ consistency error for p^ε . The value of the plateau is thus only dependent of the mesh sizes and does not depend on the values of ε . With vanishing values of ε the round-off errors due to the linear system solver grow linearly (in log scale) until they reach the consistency error ($\mathcal{O}(h^2)$). This occurs for a value of ε which, for this test case, can be estimated as approximately $\varepsilon = 10^{-9}$. For smaller ε , the discretization error is negligible compared to the round-off errors amplified by the factor $1/\varepsilon$ and the accuracy of p^ε deteriorates linearly with vanishing ε .

The accuracy of the approximation of p^ε can be made totally independent of ε under the assumption that $f^{(0)} = 0$. In this case, both $b \cdot \nabla g^\varepsilon$ and f^ε scale as ε , providing then an approximation of p^ε independent of ε . The numerical methods introduced in [10, 12] have been developed under this assumption that $f^{(0)} = 0$. The present paper is developed




Figure 1: Oblique magnetic field test case with $\alpha = \pi/3$: error norms, defined by (4.1), for the solution ϕ^ϵ as a function of the mesh size, in decimal logarithmic scales and for different values of ϵ .




Figure 2: Oblique magnetic field test case for $\alpha = \pi/3$ and $\Delta x = \Delta y = 1/60$. Approximation of the p^ϵ part of the solution.

under a weaker hypothesis, required by the application to the Euler-Lorentz model in the drift-limit. This explains why a comparable accuracy cannot be reached. Therefore, strictly speaking, our scheme is AP for the computation of p^ϵ only when $f^{(0)} = 0$, or, when




Figure 3: Oblique magnetic field test case for $\varepsilon=10^{-9}$ and $\Delta x=\Delta y=1/40$. Norms of the relative error (4.1) as a function of the angle of the magnetic field with the x -axis α .

$f^{(0)} \neq 0$, only if the round-off errors brought by the linear system solver are smaller than the discretization error. Still, it is AP without any restriction for the computation of q^ε (i.e., even when $f^{(0)} \neq 0$).

The next simulation is aimed at investigating whether the accuracy depends on the angle between b and the coordinate axes. For this purpose, simulations are carried out on a mesh composed of 40×40 cells and for α ranging from 0 to $\pi/2$. When $\alpha=0$ the b field is aligned with the x -axis and when $\alpha=\pi/2$, it is aligned with the y -axis. The relative errors are displayed as functions of α on Fig. 3. We observe that the variations of the errors are small on the whole range of angles. This confirms that the method provides accurate results, even when the mesh is far from consistent with the b -field direction.

4.2.3 Inhomogeneous Neumann boundary conditions

We remark that $\phi^\varepsilon(x,y) = 2x^2 + y^2$ is an analytical solution of system (2.28a), (2.28b) for $f_2(x,y) = \varepsilon(2x^2 + y^2)$ and $\kappa = -\nabla \cdot (bf)$. For this analytical solution and $\varepsilon=10^{-9}$, we have checked that the relative error does not exceed 10^{-13} .

4.3 Numerical results for a non uniform magnetic field

4.3.1 Introduction and test case settings

In this subsection $\Omega = [1,2] \times [1,2]$ and the magnetic field is given by:


$$B = |B|b, \quad b = (\sin(\theta), -\cos(\theta)), \quad \tan(\theta) = \frac{y}{x}. \quad (4.3)$$

For this case, an analytical solution of (1.1a), (1.1b) can be found. We consider H_{var} defined on $[1,2] \times [1,2]$ by $H_{var}(x,y) = (1-x)^3(1-y)^3(2-x)^3(2-y)^3$. According to Theorem 2.1, $b \cdot \nabla H_{var} \in K^\perp$. So $\phi = 1 + b \cdot \nabla H_{var}$ is the solution of (1.1a), (1.1b) when the right-hand f^ε of (1.1a) has the expression


$$f^\varepsilon = -b \cdot \nabla (\nabla \cdot (b \otimes b) \nabla H_{var}) + \varepsilon (1 + b \cdot \nabla H_{var}).$$

4.3.2 Homogeneous Neumann boundary conditions


On Fig. 4(a), Fig. 4(b) and Fig. 4(c), we have represented the relative errors as functions of the mesh size when ε goes from 10^{-3} to 10^{-9} . We observe that all the three norms decrease when the mesh sizes decrease, in a similar fashion as in the oblique uniform b -field.



(a) $\varepsilon = 10^{-3}$



(b) $\varepsilon = 10^{-6}$



(c) $\varepsilon = 10^{-9}$

Figure 4: Non uniform magnetic field test case: error norms, defined by (4.1), for the solution ϕ^ε as a function of the mesh size, in decimal logarithmic scales and for different values of ε .

4.3.3 Inhomogeneous Neumann boundary conditions

We take the test case of Subsubsection 4.2.3 again and we find a similar conclusion: with $\varepsilon = 10^{-9}$, the relative error in L^∞ norm does not exceed 10^{-11} .

5 Application to the Euler-Lorentz system in the drift limit

5.1 Introduction

In this section the drift-fluid limit of the isothermal Euler-Lorentz is investigated. This regime is representative of strongly magnetized plasma, for which the pressure term equilibrates the Lorentz force. It is obtained by letting a dimensionless parameter ε , representing the non-dimensional gyro-period as well as the square Mach number, go to zero. This limit is singular because the momentum equation in the direction of the magnetic field degenerates. Since the field may not be uniformly large, we wish to derive an Asymptotic-Preserving (AP) scheme which guarantees accurate discretizations of both the limit regime for strongly magnetized plasma ($\varepsilon \ll 1$) and the standard Euler-Lorentz system when the field strength is mild ($\varepsilon \sim 1$). With this aim, the Euler-Lorentz system is discretized in time by a semi-implicit scheme.

This scheme has already been studied in [13] for a uniform and constant magnetic field aligned with one coordinate and for physically less meaningful Dirichlet boundary conditions. The present methodology allows us to investigate the case of non-uniform magnetic fields and Neumann boundary conditions. Indeed, the anisotropic elliptic equation (1.1a), (1.1b) appears as the central building block of the scheme, which allows for the computation of the field-aligned momentum component. In this presentation, we will mainly focus on this aspect, the other ones being unchanged compared to [13].

5.2 The Euler-Lorentz model and its drift-fluid limit

5.2.1 The Euler-Lorentz model

The scaled isothermal Euler-Lorentz model takes the form:

$$\partial_t n_\varepsilon + \nabla \cdot (n_\varepsilon u_\varepsilon) = 0, \quad (5.1a)$$

$$\varepsilon [\partial_t (n_\varepsilon u_\varepsilon) + \nabla \cdot (n_\varepsilon u_\varepsilon \otimes u_\varepsilon)] + T \nabla n_\varepsilon = n_\varepsilon (E + u_\varepsilon \times B), \quad (5.1b)$$

where n_ε , u_ε and T are the density, the velocity and the temperature of the ions, respectively. Here, the electric field E and the magnetic field B are assumed to be given functions. The parameter ε is related to the gyro-period of the particles about the magnetic field lines and simultaneously to the squared Mach number. We refer to [13] for more details on the model, the scaling and the drift-fluid limit $\varepsilon \rightarrow 0$ but for the sake of clarity we recall some elements up to the end of this subsection.

5.2.2 The drift-fluid limit

The formal limit $\varepsilon \rightarrow 0$ in the isothermal Euler-Lorentz model (5.1a), (5.1b), leads to the so-called isothermal drift-fluid model:

$$\partial_t n + \nabla \cdot (nu) = 0, \quad (5.2a)$$

$$T \nabla n = n(E + u \times B). \quad (5.2b)$$

The constraint (5.2b) completely determines the velocity \mathbf{u} . Indeed, taking the parallel and perpendicular components of (5.2b) leads to

$$nu_{\perp} = \frac{1}{B} b \times (T \nabla n - nE), \quad (5.3a)$$

$$T \nabla_{\parallel} n - nE_{\parallel} = 0. \quad (5.3b)$$

After dividing by n , we find that the first term at the right-hand side of (5.3a) is the diamagnetic drift velocity while the second one is the $E \times B$ drift velocity. Note that, by arguing as in [13] Eq. (5.3b) can be written in the form of an elliptic equation for nu_{\parallel} as follows:

$$-\nabla_{\parallel} (\nabla_{\parallel} \cdot (nu_{\parallel})) = \partial_t \left(\frac{nE_{\parallel}}{T} \right) - \partial_t b \cdot \nabla n + \nabla_{\parallel} \nabla_{\perp} \cdot (nu_{\perp}). \quad (5.4)$$

The drift-fluid model consists of Eqs. (5.2a), (5.3a) and (5.4).

5.2.3 A reformulation of the isothermal Euler-Lorentz model

As recalled in the introduction of this section, to find an AP scheme, it is essential to "regularize" the perturbation, i.e., to reformulate the Euler-Lorentz model in such a way that the limit equations for the velocity appear explicitly in the system of equations. By reasoning as in [13] we get that nu_{\perp} and nu_{\parallel} satisfy

$$\begin{aligned} & B(nu)_{\perp} - \varepsilon \partial_t (b \times (nu)_{\perp}) \\ &= -b \times [-T \nabla n + nE] + \varepsilon [-(\partial_t b) \times (nu) + b \times (\nabla \cdot (nu \otimes u))] \end{aligned} \quad (5.5a)$$

and

$$\begin{aligned} & \varepsilon \partial_t^2 ((nu_{\varepsilon})_{\parallel}) - T \nabla_{\parallel} (\nabla_{\parallel} \cdot (nu_{\varepsilon})_{\parallel}) \\ &= \varepsilon \partial_t ((\partial_t b) \cdot ((nu)_{\parallel})) - \varepsilon \partial_t (b \cdot (\nabla \cdot ((nu_{\varepsilon} \otimes u_{\varepsilon}))) + \partial_t (n_{\varepsilon} E_{\parallel}) \\ & \quad - T \partial_t b \cdot \nabla n_{\varepsilon} + T \nabla_{\parallel} (\nabla_{\perp} \cdot (n_{\varepsilon} u_{\varepsilon})_{\perp}). \end{aligned} \quad (5.5b)$$

Therefore, the reformulation of the Euler-Lorentz model consists of Eqs. (5.1a), (5.5a) and (5.5b).

5.3 The AP scheme

Now we introduce the time discretization of the model. Let B^m be the magnetic field at time t^m , $|B|^m$ its magnitude and $b^m = B^m / |B|^m$ its direction. For a given vector field v , denote by $(v)_{\parallel}^m$ and $(v)_{\perp}^m$ its parallel and perpendicular components with respect to b^m i.e.,

$$v = (v)_{\parallel}^m b^m + (v)_{\perp}^m, \quad (v)_{\parallel}^m = v \cdot b^m, \quad (v)_{\perp}^m = b^m \times (v \times b^m).$$

Similarly, we denote by ∇_{\parallel}^m and $\nabla_{\perp}^m \cdot$ the parallel gradient and divergence operators respective to this field. The time semi-discrete scheme proposed in [13] is as follows:

Definition 5.1. The AP scheme is the scheme defined by:

$$\frac{n^{m+1} - n^m}{\Delta t} + \nabla \cdot (nu)^{m+1} = 0, \quad (5.6a)$$

$$\varepsilon \left[\frac{(nu)^{m+1} - (nu)^m}{\Delta t} + \nabla \cdot (nu \otimes u)^m \right] + T(\nabla n^\#)^{m+1} = n^m E^{m+1} + (nu)^{m+1} \times B^{m+1}, \quad (5.6b)$$

where $(\nabla n^\#)^{m+1}$ is given by

$$(\nabla n^\#)^{m+1} = (\nabla n^m)_\perp^{m+1} + (\nabla n^{m+1})_\parallel^{m+1} b^{m+1}. \quad (5.7)$$

The cross-product of (5.6b) with b^{m+1} leads to

$$\begin{aligned} & (nu)_\perp^{m+1} - \frac{\varepsilon}{\Delta t} \frac{1}{B^{m+1}} b^{m+1} \times (nu)_\perp^{m+1} \\ &= -\frac{1}{B^{m+1}} b^{m+1} \times \left[\frac{\varepsilon}{\Delta t} (nu)^m - \varepsilon \nabla \cdot (nu \otimes u)^m - T \nabla n^m + n^m E^{m+1} \right], \end{aligned} \quad (5.8)$$

which is a discretization of Eq. (5.5a), where

$$(\partial_t b) \times (nu) \approx \left(\frac{b^{m+1} - b^m}{\Delta t} \right) \times (nu)^m.$$

By considering the scalar product of (5.6b) with b^{m+1} , we get

$$\varepsilon \left(\frac{(nu)^{m+1} - (nu)^m}{\Delta t} + \nabla \cdot (n_\varepsilon u_\varepsilon \otimes u_\varepsilon)^m \right) \cdot b^{m+1} = -T \nabla^{m+1} n^m \cdot b^{m+1} + n^m E^{m+1} \cdot b^{m+1}$$

and after easy computations [13], we find that $(nu)_\parallel^{m+1}$ satisfies the following anisotropic elliptic problem:

$$\begin{aligned} & \frac{\varepsilon}{\Delta t} (nu)_\parallel^{m+1} - T \Delta t \nabla_\parallel^{m+1} \left(\nabla_\parallel^{m+1} \cdot ((nu)^{m+1})_\parallel^{m+1} \right) \\ &= T \Delta t \nabla_\parallel^{m+1} \left(\nabla \cdot ((nu)^{m+1})_\perp^{m+1} \right) - T \nabla_\parallel^{m+1} n^m \\ &+ \left[\frac{\varepsilon}{\Delta t} (nu)^m - \varepsilon (\nabla \cdot (nu \otimes u)^m) + n^m E^{m+1} \right]_\parallel^{m+1}. \end{aligned} \quad (5.9)$$

By setting $(nu)_\parallel^{m+1} = \phi^\varepsilon$ and by taking $f = f_1 + f_2$ with

$$f_1 = \frac{1}{\Delta t} b \cdot \nabla (\nabla \cdot (nu)_\perp^{m+1}), \quad (5.10a)$$

$$f_2 = - \left[\frac{\varepsilon}{T(\Delta t)^2} (nu)^m - \frac{\varepsilon}{T \Delta t} \nabla \cdot (nu \otimes u)^m + n^m E^{m+1} \right]_\parallel^{m+1} - \frac{1}{\Delta t} (b \cdot \nabla n^m), \quad (5.10b)$$

this problem can be put in the framework of (1.1a). In [13], because b was chosen parallel to one of the coordinate axes, a direct discretization of (5.9) using finite differences could

be performed. Here, for an arbitrary anisotropy direction b , we use the method developed in the previous sections. We do not detail the description of the discretization of the other equations, since it follows [13].

The right-hand side (5.10b) can be decomposed as $f_2^\varepsilon = f_2^{(0)} + \varepsilon f_2^{(1)}$ with $f_2^{(0)}$ corresponding to the first two terms and $f_2^{(1)}$, to the last two one. Moreover if we suppose that

$$[n^m E^{m+1}]_{\parallel}^{m+1} - \frac{1}{\Delta t} (b \cdot \nabla n^m) \in K^\perp, \quad (5.11)$$

the compatibility condition (2.25) is satisfied. This property amounts to saying that the integrated force along a magnetic field line is zero. If the property is not satisfied, parallel velocities of order $\mathcal{O}(\varepsilon^{-1})$ are generated, which is physically unrealistic (because collisions will ultimately slow down the plasma ions). Therefore, assuming (5.11) is physically justified.

Next for the sake of clarity we rewrite the Eqs. (5.8), (5.9) in the situation where $b = (b_x, b_y, 0)$. In that case

$$\nabla_{\parallel}^{m+1} (\nabla \cdot (nu_\varepsilon)_\perp^{m+1})^{m+1} = b_x \partial_x (\partial_x (b_y^2 nu_x - b_y b_x nu_y)) + b_y \partial_y (\partial_y (b_x^2 nu_y - b_y b_x nu_x))$$

and

$$\nabla \cdot (nu^\varepsilon \otimes u^\varepsilon) \cdot b = b_x (\partial_x (nu_x^2) + \partial_y (nu_x u_y)) + b_y (\partial_x (nu_x u_y) + \partial_y (nu_y^2)).$$

Therefore the parallel velocity satisfies

$$\begin{aligned} & \frac{\varepsilon}{\Delta t} (nu)_{\parallel}^{m+1} - T \Delta t \nabla_{\parallel}^{m+1} (\nabla_{\parallel}^{m+1} \cdot ((nu)^{m+1})_{\parallel}^{m+1}) \\ &= T (b_x \partial_x + b_y \partial_y) n^m + b_x (\partial_x (nu_x^2)^m + \partial_y (nu_x u_y)^m) \\ & \quad + b_y (\partial_x ((nu_x u_y)^m) + \partial_y ((nu_y^2)^m)) + \varepsilon b_x \partial_x (\partial_x (b_y^2 (nu_x)^{m+1} - b_y b_x (nu_y)^{m+1})) \\ & \quad + \varepsilon b_y \partial_y (\partial_y (b_x^2 (nu_y)^{m+1} - b_y b_x (nu_x)^{m+1})) + n^m E_{\parallel}^{m+1} + \frac{\varepsilon}{\Delta t} (nu)_{\parallel}^m. \end{aligned} \quad (5.12)$$

Recall that the orthogonal part of the velocity is obtained from (5.8). So in the present case it is determined by inverting the system

$$(nu)_{\perp}^{m+1} - \frac{\varepsilon}{\Delta t} \frac{1}{B^{m+1}} b^{m+1} \times (nu)_{\perp}^{m+1} = (V_1, V_2, V_3),$$

where

$$\begin{aligned} V_1 &= b_y \left(\frac{\varepsilon}{\Delta t} (nu_z)^m - n^m E_z^{m+1} \right), & V_2 &= b_x \left(\frac{\varepsilon}{\Delta t} (nu_y)^m + n^m E_z^{m+1} \right), \\ V_3 &= b_x \left(\frac{\varepsilon}{\Delta t} (nu_y)^m + n^m E_y^{m+1} + \varepsilon (\partial_x (nu_x u_y) + \partial_y (nu_y^2)) - T \partial_y n \right) \\ & \quad - b_y \left(\frac{\varepsilon}{\Delta t} (nu_x)^m + n^m E_x^{m+1} + \varepsilon (\partial_y (nu_x u_y) + \partial_x (nu_x^2)) - T \partial_x n \right). \end{aligned}$$

As in [13], we will compare the AP scheme with the classical semi-discrete scheme for the Euler-Lorentz model, given by:

Definition 5.2. The "classical" semi-discrete scheme is defined by:

$$\frac{n^{m+1} - n^m}{\Delta t} + \nabla \cdot (nu)^m = 0, \quad (5.13a)$$

$$\varepsilon \left[\frac{(nu)^{m+1} - (nu)^m}{\Delta t} + \nabla \cdot (nu \otimes u)^m \right] + T(\nabla n)^m = n^m E^{m+1} + (nu)^{m+1} \times B^{m+1}. \quad (5.13b)$$

In [13], it is shown that this scheme is not uniformly stable with respect to ε and so that it cannot be AP.

Except from the parallel momentum equation, which has just been discussed, the other equations of the model are discretized following [13]. For the sake of brevity, we will not reproduce their presentation here.

5.4 Boundary conditions

The AP-scheme involves the resolution an elliptic equation for u_{\parallel} which requires boundary conditions. These boundary conditions should in principle derive from the original formulation of the Euler-Lorentz model. However, there is no obvious way to derive them explicitly in the general case and in practice, they must be set up on a case by case basis, from the knowledge of the underlying physical situation. In this paper, our goal is to show that the AP-scheme is stable with under-resolved time steps, i.e., time steps which are large compared to the times scales linked to ε . For this purpose, we propose boundary conditions which strictly speaking are valid only in the $\varepsilon \rightarrow 0$ limit. These approximate boundary conditions are the reason for some small discrepancies with the true solution near the domain boundaries as we will see on the numerical results. Still, we will observe that the AP-scheme shows the desired stability property.

We impose Dirichlet boundary conditions on the density $n^{m+1} = n_B$ with n_B independent of time. For the perpendicular momentum, we impose the relation obtained after taking the limit when $\varepsilon \rightarrow 0$ in (5.1b),

$$nu_{\perp}^{m+1} = -\frac{1}{|B|^{m+1}} b \times (T\nabla n^m + n^m E^{m+1}).$$

By considering the mass conservation equation at the domain boundary, we have

$$\frac{n^{m+1} - n^m}{\Delta t} + \nabla \cdot (bn^{m+1}u_{\parallel}^{m+1}) + \nabla \cdot (nu)_{\perp}^{m+1} = 0, \quad \text{on } \partial\Omega.$$

Therefore, as the density satisfies Dirichlet boundary conditions with time-independent Dirichlet values, we get

$$(b \cdot \nu) \nabla \cdot (bnu_{\parallel}^{m+1}) = -(b \cdot \nu) \nabla \cdot (nu_{\perp}^{m+1}), \quad \text{on } \partial\Omega.$$

Therefore, nu_{\parallel}^{m+1} is a solution to the anisotropic elliptic problem with inhomogeneous Neumann boundary conditions (2.28a), (2.28b), with $\kappa = -(b \cdot \nu) \nabla \cdot (nu_{\perp}^{m+1})$. Then, we can

apply the framework of Section 2.2.3. When nu_{\parallel} has been calculated, an approximation is employed in order to provide values of nu_{\parallel} in a layer of fictitious cells surrounding the boundary, by using homogeneous Neumann boundary conditions. The values in the fictitious cells are then useful to compute gradient terms which occur in the other equations of the Euler-Lorentz model.

6 Numerical results for the Euler-Lorentz system in the drift limit

6.1 Introduction and test case settings

This part is devoted to the validation of the AP-scheme (5.6a), (5.6b), (5.7) for the Euler-Lorentz system. Due to the lack of analytical solutions, the validation procedure consists in comparisons of the AP-scheme with the classical discretization (5.13b). The classical discretization is subject to a CFL stability condition that imposes the time step to resolve (i.e., to be smaller than) the fastest time scales involved in the system. These *time-resolved* simulations require a time step which scales like $\sqrt{\varepsilon}$ (because the CFL condition involves the acoustic wave speed which scales like $1/\sqrt{\varepsilon}$). The AP-scheme is designed to be stable independently of ε when $\varepsilon \rightarrow 0$. In these situations, the time step cannot resolve the fastest time scales involved in the system, which leads to *under-resolved* simulations. The stability of the AP-scheme in under-resolved situations has been demonstrated in [13]. In this case, the requested CFL condition only involves the fluid velocity, which is an $\mathcal{O}(1)$ quantity and not the acoustic speed [13] and explains the possibility of using large time steps, independent of ε . We want to check this feature again when the scheme is equipped with our new elliptic solver.

Two test cases are presented, one for an oblique uniform magnetic field, another one for a non uniform magnetic field with the same expressions as in Section 4. In both cases, the electric field is chosen as $E = (0, 0, B_x + B_y)$, where B_x and B_y are the components of the magnetic field. The initial condition is defined by the following uniform data: $n = 1$, $(nu)_x = 1$, $(nu)_y = -1$ and $(nu)_z = 0$ which defines a stationary solution of the Euler-Lorentz system. A local perturbation of order ε is then applied to this stationary state and the evolution of the system is observed for both the AP and the classical schemes.

6.2 Numerical results for an oblique uniform magnetic field

The results for the AP and the classical schemes are compared on Fig. 5 in a resolved case. Both schemes provide comparable results. However we observe the formation of a thin boundary layer on the domain frontiers for the AP-scheme but it is not responsible for the development of an instability. The occurrence of this boundary layer is due to the additional boundary condition, as explained in Section 5.4.

Next we consider the same test case with an under-resolved time step Δt which is 10 times larger than the time step provided by the CFL condition of the classical scheme.







(a) n (AP-scheme)(b) n (classical scheme)(c) nu_x (AP-scheme)(d) nu_x (classical scheme)(e) nu_y (AP-scheme)(f) nu_y (classical scheme)

Figure 5: Euler-Lorentz test case for an oblique magnetic field in the resolved case at time $t=3.9510^{-6}$ s: density (n) and momentum (nu_x , nu_y) computed by the AP-scheme (left) and the classical scheme (right) for $\varepsilon=10^{-9}$ and $\Delta x=\Delta y=1/40$. The angle of the magnetic field with the x -axis is $\alpha=\pi/3$.

These simulation results are collected on Fig. 6. The results are displayed at the time 3.9510^{-5} s, which is 10 times bigger than the time displayed in Figs. 5(a)-5(f). This larger simulation time is used to show the capability of the AP scheme to sustain long simulation times. In this case, the conventional scheme leads to unstable results contrary to the AP scheme and proves the capability of the AP-scheme to provide stable computations for time steps that resolve neither the acoustic wave-speed nor the gyration period. We







(a) n (AP-scheme)(b) n (classical-scheme)(c) nu_x (AP-scheme)(d) nu_x (classical-scheme)(e) nu_y (AP-scheme)(f) nu_y (classical-scheme)

Figure 6: Euler-Lorentz test case for an oblique magnetic field in the under-resolved case at time 3.9510^{-5} s: density (n) and momentum (nu_x , nu_y) computed by the AP-scheme (left) and the classical scheme (right) for $\varepsilon = 10^{-9}$ and $\Delta x = \Delta y = 1/40$. The angle of the magnetic field with the x -axis is $\alpha = \pi/3$.

also note that, over these simulation times, the AP-scheme produces the same results in the resolved and under-resolved cases (not displayed). In the numerical test, the perturbation is localized in the center of the domain at the beginning of the simulation and spreads out until it reaches the boundaries of the domain. In Figs. 6(a)-6(f), we observe the solution after the perturbation has reached the boundary of the domain, while in

Figs. 5(a)-5(f), we observe it at a shorter time when it is still well inside the domain. That is why the solutions obtained by the AP scheme on the two figures look different.

6.3 Numerical results for a non uniform magnetic field

For the non uniform case, $n = 1$, $(nu)_x = 1$, $(nu)_y = -1$ and $(nu)_z = 0$ are not stationary solutions to the Euler-Lorentz system. In particular, with the chosen initial condition,




Figure 7: Euler-Lorentz test case for a non uniform magnetic field in the resolved case at time $t = 3.9510^{-6}$ s: density (n) and momentum (nu_x , nu_y) computed by the AP-scheme (left) and the classical scheme (right) for $\epsilon = 10^{-9}$ and $\Delta x = \Delta y = 1/40$.







(a) n (AP-scheme)(b) n (classical-scheme)(c) nu_x (AP-scheme)(d) nu_x (classical-scheme)(e) nu_y (AP-scheme)(f) nu_y (classical-scheme)

Figure 8: Euler-Lorentz test case for a non uniform magnetic field in the under-resolved case at time $t=3.9510^{-5}$ s: density (n) and momentum (nu_x , nu_y) computed by the AP-scheme (left) and the classical scheme (right) for $\varepsilon=10^{-9}$ and $\Delta x=\Delta y=1/40$.

sharp boundary layers are generated. But the AP scheme can still be compared with the classical scheme in the resolved case for a validation procedure. Then we take the same initial conditions as for the oblique magnetic field case. Fig. 7 shows that the two schemes provide similar results.

Next we consider the under-resolved time step $10\Delta t$. In this situation Figs. 8(b), 8(d),




(a) Difference for n (b) Difference for nu_x (c) Difference for nu_y

Figure 9: Difference between the solutions obtained with an initial perturbation of order $\varepsilon = 10^{-9}$ and the solution without any perturbation for the variable magnetic field after 1.58s of simulation in the non resolved case.

8(f) show that the classical scheme is unstable. By contrast, Figs. 8(a), 8(c), 8(e) demonstrate that the AP-scheme provides stable results. The increased numerical diffusion generated by the large time step gives rise to a widening of the boundary layer. Keeping the boundary layer accurate would require some mesh refinement in the vicinity of the boundary. This point is deferred to future work.

Moreover as the initial conditions of the present test case are not stationary solutions of the Euler-Lorentz model, it is important to check if the results obtained in the non resolved case by the AP scheme correspond to the proper limit regime. So we compare the results obtained with and without the local perturbation on the initial conditions. The difference between the results obtained with the two simulations remain of the same order as the perturbation of the initial condition. Figs. 9(a), 9(b) and 9(c) present the difference between the solutions obtained with the perturbed and non-perturbed initial condition, for n , $(nu)_x$, $(nu)_y$ after a very long simulation time of 1.58s, about 10^5 longer than in Figs. 8(a)-8(f). The figures show that this difference is actually of 10^{-10} for the density and 10^{-6} for the momenta. The difference with the value of $\varepsilon = 10^{-9}$, can be explained by the accumulation of the truncation error over the very large simulation time.

7 Conclusions and perspectives

A numerical method for degenerate anisotropic elliptic problems has been investigated. This method is based on a variational formulation together with a decomposition of the solution. This problem has been applied to the resolution of an Asymptotic-Preserving scheme for the isothermal Euler-Lorentz system. Numerical simulations demonstrate the ability of the scheme to handle under-resolved situations where the time-step exceeds the CFL stability condition of the classical scheme.

Forthcoming works will be devoted to the generalization of this approach for the full Euler system with a non linear pressure law. In this case non linear anisotropic elliptic problem have to be handled. Moreover we can also deal with the more physical situation of a plasma constituted by a mixture of ions and electrons. In this situation the model can be described by the two-fluid Euler-Lorentz system coupled with quasi-neutrality equation.

Acknowledgments

This work has been partially supported by the Marie Curie Actions of the European Commission in the frame of the DEASE project (MEST-CT-2005-021122), by the "Fédération de recherche CNRS sur la fusion par confinement magnétique", by the Association Euratom-CEA in the framework of the contract "Gyro-AP" (contract # V3629.001 avenant 1) and by the University Paul Sabatier in the frame of the contract "MOSITER". This work was performed while the first author held a post-doctoral position funded by the Fondation "Sciences et Technologies pour l'Aéronautique et l'Espace", in the frame of the project "Plasmax" (contract # RTRA-STAE/2007/PF/002). The authors wish to express their gratitude to G. Falchetto, X. Garbet and M. Ottaviani from CEA-Cadarache and G. Gallice and C. Tessieras from CEA-Cesta for fruitful discussions and encouragements.

References

- [1] M. Beer, S. Cowley and G. Hammett, Field-aligned coordinates for nonlinear simulations of tokamak turbulence, *Phys. Plasmas*, 2(7) (1995), 2687–2700.
- [2] M. A. Beer and G. W. Hammett, Toroidal gyrofluid equations for simulations of tokamak turbulence, *Phys. Plasmas*, 3 (1996), 4046–4064.
- [3] R. Belaouar, N. Crouseilles, P. Degond and E. Sonnendrcker, An asymptotically stable semi-lagrangian scheme in the quasi-neutral limit, *J. Sci. Comput.*, (2009), 341–365.
- [4] C. Besse, F. Deluzet, C. Negulescu and C. Yang, Three dimensional simulation of ionospheric plasma disturbances, in preparation.
- [5] A. H. Boozer, Establishment of magnetic coordinates for a given magnetic field, *Phys. Fluids*, 25(3) (1982), 520–521.
- [6] C. Buet, S. Cordier, B. Lucquin-Desreux and S. Mancini, Diffusion limit of the Lorentz model: Asymptotic Preserving schemes, *ESAIM: M2AN*, 36 (2002), 631–655.

- [7] C. Buet and B. Despres, Asymptotic Preserving and positive schemes for radiation hydrodynamics, *J. Comput. Phys.*, 215 (2006), 717–740.
- [8] J. A. Carrillo, T. Goudon and P. Lafitte, Simulation of fluid and particles flows: Asymptotic Preserving schemes for bubbling and flowing regimes, *J. Comput. Phys.*, 227 (2008), 7929–7951.
- [9] P. Crispel, P. Degond and M.-H. Vignal, An Asymptotic Preserving scheme for the two-fluid Euler-Poisson model in the quasineutral limit, *J. Comput. Phys.*, 223(1) (2007), 208–234.
- [10] P. Degond, F. Deluzet, A. Lozinski, J. Narski and C. Negulescu, Duality-based Asymptotic-Preserving method for highly anisotropic diffusion equations, arXiv:1008.3405v1, 2010.
- [11] P. Degond, F. Deluzet, L. Navoret, A.-B. Sun and M.-H. Vignal, Asymptotic-Preserving Particle-In-Cell method for the Vlasov-Poisson system near quasineutrality, *J. Comput. Phys.*, 229(16) (2010), 5630–5652.
- [12] P. Degond, F. Deluzet and C. Negulescu, An Asymptotic Preserving scheme for strongly anisotropic elliptic problems, *Multiscale Model. Sim.*, 8(2) (2010), 645–666.
- [13] P. Degond, F. Deluzet, A. Sangam and M.-H. Vignal, An Asymptotic-Preserving scheme for the Euler equations in a strong magnetic field, *J. Comput. Phys.*, 228(10) (2009), 3540–3558.
- [14] P. Degond, H. Liu, D. Savelief and M. H. Vignal, Numerical approximation of the Euler-Poisson-Boltzmann model in the quasineutral limit, submitted.
- [15] P. Degond, J.-G. Liu and M.-H. Vignal, Analysis of an Asymptotic Preserving scheme for the Euler-Poisson system in the quasineutral limit, *SIAM J. Numer. Anal.*, 46(3) (2008), 1298–1322.
- [16] P. Degond and M. Tang, All speed scheme for the low Mach number limit of the isentropic Euler equation, *Commun. Comput. Phys.*, to appear.
- [17] W. D. D’haeseleer, W. N. G. Hitchon, J. D. Callen and J. L. Shohet, Flux Coordinates and Magnetic Field Structure: A Guide to a Fundamental Tool of Plasma Theory, Springer Series in Computational Physics, Springer-Verlag, Berlin, 1991.
- [18] A. M. Dimits, Fluid simulations of tokamak turbulence in quasiballoon coordinates, *Phys. Rev. E*, 48(5) (1993), 4070–4079.
- [19] W. Dorland and G. Hammett, Gyrofluid turbulence models with kinetic effects, *Phys. Fluids B*, 5(3) (1993), 812–835.
- [20] F. Filbet and S. Jin, A class of Asymptotic Preserving schemes for kinetic equations and related problems with stiff sources, *J. Comput. Phys.*, in press.
- [21] L. Giraud and R. S. Tuminaro, Schur complement preconditioners for anisotropic problems, *IMA J. Numer. Anal.*, 19(1) (1999), 1–18.
- [22] V. Grandgirard, Y. Sarazin, X. Garbet, G. Dif-Pradalier, P. Ghendrih, N. Crouseilles, G. Latu, E. Sonnendrücker, N. Besse and P. Bertrand, Computing ITG turbulence with a full- f semi-lagrangian code, *Commun. Nonlinear Sci. Numer. Simulat.*, 13(1) (2008), 81–87.
- [23] S. Hamada, Hydromagnetic equilibria and their proper coordinates, *Nucl. Fusion*, 2 (1962), 23–37.
- [24] G. W. Hammett, M. A. Beer, W. Dorland, S. C. Cowley and S. A. Smith, Developments in the gyrofluid approach to tokamak turbulence simulations, *Plasma Phys. Contr. Fusion*, 35(8) (1993), 973.
- [25] S. Jin, Efficient Asymptotic-Preserving (AP) schemes for some multiscale kinetic equations, *SIAM J. Sci. Comput.*, 21(2) (1999), 441–454.
- [26] E. Kaveeva and V. Rozhansky, Poloidal and toroidal flows in tokamak plasma near magnetic islands, *Tech. Phys. Lett.*, 30(7) (2004), 538–540.
- [27] B. N. Khoromskij and G. Wittum, Robust Schur complement method for strongly anisotropic

- elliptic equations, *Numer. Linear Algebra Appl.*, 6(8) (1999), 621–653.
- [28] A. Klar, An Asymptotic Preserving numerical scheme for kinetic equations in the low Mach number limit, *SIAM J. Numer. Anal.*, 36 (1999), 1507–1527.
 - [29] M. Lemou and L. Mieussens, A new Asymptotic Preserving scheme based on micro-macro formulation for linear kinetic equations in the diffusion limit, *SIAM J. Sci. Comput.*, 31 (2008), 334–368.
 - [30] R. J. Leveque, *Finite Volume Method for Hyperbolic Problems*, Cambridge Texts in Applied Mathematics, Cambridge University Press, Cambridge, 2002.
 - [31] I. Llorente and N. Melson, Robust multigrid smoothers for three dimensional elliptic equations with strong anisotropies, ICASE Technical Report: TR-98-37, 1998.
 - [32] R. G. McClarren and B. Lowrie, The effects of slope limiting on Asymptotic-Preserving numerical methods for hyperbolic conservation laws, *J. Comput. Phys.*, 227 (2008), 9711–9726.
 - [33] J. M. Melenk, *hp*-finite element methods for singular perturbations, Volume 1796 of Lecture Notes in Mathematics, Springer-Verlag, Berlin, 2002.
 - [34] K. Miyamoto, *Controlled Fusion and Plasma Physics*, Chapman & Hall, 2007.
 - [35] M. A. Ottaviani, An alternative approach to field-aligned coordinates for plasma turbulence simulations, arXiv:1002.0748, 2010.
 - [36] D. Stern, Geomagnetic Euler potentials, *J. Geophys. Res.*, 72(15) (1967), 3995–4005.