

- [1] **(Journal Articles)** M. J. Berger and P. Collela, Local adaptive mesh refinement for shock hydrodynamics, *J. Comput. Phys.*, 82 (1989), 62-84.
- [2] **(Article accepted)** Z. J. Tan, T. Tang and Z. R. Zhang, A simple moving mesh method for one- and two-dimensional phase-field equations, *J. Comput. Appl. Math.*, to appear.
- [3] **(Preprints)** S. McKechnie, J. M. Frost, D. Pashov, P. Azarhoosh, A. Walsh, M. Schilfgaard, arXiv preprint 2017, DOI: 10.48550/arXiv.1711.00533.
- [4] **(Books without editor)** C. de Boor, Good Approximation By Splines With Variable Knots II, in Springer Lecture Notes Series 363, Springer-Verlag, Berlin, 1973.
- [5] **(Books with editor)** T. D. Tullius in *Comprehensive Supramolecular Chemistry*, Vol. 5 (Eds.: J. L. Atwood, J. E. D. Davies, D. D. MacNicol, F. Vögtle, K. S. Suslick), Pergamon, Oxford, 1996, pp. 317–343.
- [6] **(Data citations)** [dataset] Authors, Year, *Dataset title*, Data repository or archive, Version (if any), Persistent identifier (e.g. DOI).
- [7] **(Programs)** G. M. Sheldrick, SHELXS-96, Program for the Solution of Crystal Structures, University of Göttingen, Göttingen (Germany), 1996.
- [8] **(Thesis)** A. Name, Title of PhD thesis, University of Newcastle (UK), 1991.